



INVESTIGATION OF PROCESSED *PAULETIA MONANDRA* (KURZ) SEED-MEAL ON THE GROWTH AND COST VIABILITY OF *CLARIS GARIEPINUS* FINGERLINGS (TEUGELS) FED VARYING LEVEL OF INCLUSION IN SUSTAINABLE FISH PRODUCTION IN FEDERAL COLLEGE OF EDUCATION, TECHNICAL BICHI, NIGERIA

MAMUDU SAMSON TOYOSI and DARDA FOLORENCE Ph.D

Dept of Biology Education, Federal College of Education (T) Bichi, Kano.
Dept Integrated Sc Education, Federal College of Education (T) Bichi, Kano.

Abstract

Pauletia monandra meal was experimented for its ability and inclusion to partially replace fishmeal in the diet of *Claria gariepinus* fingerlings under out-doors culture system. The study was conducted in Biological garden, fisheries unites department of Biology Education, Federal college of Education (tech) Bichi. Kano State, Nigeria (8-9°E and 12-13°N) Five Feeds were formulated to meet the crude protein value of the control diet which is top feeds at 49.19%. The diets were fed to fish at 5% body weight thrice per day (8-9 am, 1-2pm and 4-5 pm). The experiment was conducted for 54 days, measured among other parameters are feed conversion ratio (FCR), specific growth rate (SGR) stander length (S) percentage survival rate of 88% was achieved. There was no significant difference ($P>0.05$) in the growth parameters and the control diet in the fish samples. Although *P. monandra* supported the growth performance and feed utilization was favored by low inclusion. Diet 1 was the most economical in terms of its relative cost of feed per unit weight gain of fish with only N7.09 diet per gram of weight gain, diet 5 (topfeeds) was the most expensive (N170.00 kg). The combination of *P. monandra* seeds meals formed a cheaper and better balanced diet for fingerlings of *C gariepinus* the use of processed *P. monandra* seeds meals to supplement artificial feed in catfish production is cost effective and has high potential.

Key Words: Growth, Performance, Survival, Standard Length Fingerlings Concrete Pond



This research was sponsored by Tertiary Education Trust Fund (TETFUND) research grant award through Federal College of Education (Technical) Bichi, Kano

Introduction

Aquaculture shares the same challenges with Agriculture in increasing food supply and this brings about competition in the use of feeds for livestock and fish farming. Shortages of major feedstuff has been on the increase in recent times in Nigeria and with the poultry and livestock industry expanding, the aquaculture industry is finding it increasingly more difficult to source for critical feed ingredients (Adewola 2008, Agbugui, Oniye and Auta, 2010, Mamudu, Aminu and Hassan, 2018). For aquaculture to supply the population's growing demand for fish as food and to fill the gap in declining yield from capture fisheries, basic but critical information should be available especially regarding feed that are less competitive and of low cost value but with replaceable capacity for fish meals with the aim of making fish to attain table size at reduced culture time and minimum production cost (Umaru, Auta, Oniye and Bolorunduro, 2016, Burke, 2019).



Utilization of non-conventional protein supplements of both animal and plants origin in practical fish diets has been a focus in Nigeria in recent times. It is imperative for such practical diet to contain optimum protein, required essential amino and fatty acids (F.A.O, 2020). Recently, the analysis of nutritional value of wild plant seeds attracted attention, they have shown to contain significant amount of essential nutrients (proteins, amino acids, vitamins, minerals, oils and carbohydrates) that can be used for the formulation of animal feeds, In view of the presence of anti-nutritional factors in these seeds, they have to be processed to deactivate the protease inhibitors, or reduced to a level that can be tolerated by the fish or completely eliminated (Anhwange, Ajibola, and Oniye, 2005, Anwa, Auta, Abdullahi and Bolorunduro, 2017 and Salgueiro, 2022). *P. monandra* is typically a savannah tree found in Nigeria. Its seeds have been subjected to a variety of uses. For example, Teixeira, Farias, Sousa and Vasconcelos, (2013) reported that diets made from *P. monandra* seeds were used to feed Japanese quails and the diet enhanced growth and the onset of early egg laying. The seeds have a high protein content which is found to be 33% (Ferouz, Kurt and Kasiuviswanathan, 2012). Various qualities such as availability (during the dry season), ease of propagation and simple processing techniques make *P. monandra* a choice for research into its suitability for compounding fish feed formulation for *C. gariepinus*. The African cat fish has been chosen for this study because of its importance in aquaculture. *C. gariepinus* is one of the culture able fish species because it is hardy, can withstand stress, its rapid growth rate, acceptability of variety of food items and its ability to respond to induced breeding, Omosowone and Ogunrinde, (2018). The fish is highly priced with high economic returns as fresh or smoked or dried fish with high demand (Abduilah, *et al* 2009, Tunde, Babatunde and Oluseyi 2018).

Objectives; to investigate the possibility of inclusion of processed *P. monandra* seed meal in diet of *C. gariepinus*

- To determine the growth performance of *C. gariepinus* fingerlings fed on diets containing processed *P. monandra* seed meal and topfeeds.
- To assess the cost of feeding *C. gariepinus* with diet containing *P. monandra* seed meal and Top feed.

Hypotheses

- There is no significant differences between the growth performance *C. gariepinus* fingerlings fed on diets containing boiled *P. monandra* seed meal and top feed meal .
- There is no significant difference in the cost of feeding *C. gariepinus* fingerlings with diets containing *P. mnandra* seed meal and top feed.



Pods



Drying



Harvesting

MATERIALS AND METHODS.

Diet formation and preparation. Freshly harvested dry pod of *P. monandra* from Biological garden of the Department of Biological Sciences Ahmadu Bello University, Zaria, Nigeria was collected, drying and package. Other feeds ingredients such as yellow maze, fish meal, cassava starch, vitamin



premix and salt, were purchased locally from market from Badume markets and commercial feed stores in Bichi, Kano. Five portions of *P. monandra seeds* (100g each) were boiled at different time intervals of 10, 20, 30, 40 and 60 minutes corresponding to T2, T3, T4, T5 and T6 respectively, the six portion T1 was raw and not boiled the boiling process was carried out using a pressure pot with about 3L of water on gas burner. The boiled seeds were then air dried for one week in the laboratory and grounded into fine powder and label and packaged in labeled polythene bags. Proximate analysis and the level of their ant- nutritional content were also determined. All dietary ingredients were milled using locally fabricated hammer mill and sieved through a 595um sieve to remove stones and dirt as well as ensure homogeneous size profile before being analyzed for proximate composition including mineral premix/ vitamin supplement and table salt were added and mixed during grinding to ensure homogeneousness. The binder was made by first making a 3g cassava paste and then adding about 2 liters of hot water to the paste while constantly stirring the past. The powered diets prepared were mixed with cassava paste to make dough and then made into pellet (2millimeters) from using a modified small size hammer mill pelletizing machine. The pellet was air dried for one week and bagged. The proximate analyses of the diets were determined by the method described by AOAC (2020).



Preparation of fish feed ingredient

Table 1. Percentage composition of Experimental Diets for *C. gariepinus* fingerlings

Ingredients (kg)	Diet 1	Diet 2	Diet 3	Diet 4	Diet 5
Fish meal	31.10	35.30	39.70	44.00	Nil
Maize	5.00	3.00	1.00	Nil	"
Palm oil	5.58	3.38	1.15	Nil	"
Cassava starch	2.00	2.00	2.00	2.00	"
Premix	0.25	0.25	0.25	0.25	"
Salt	0.30	0.30	0.30	0.30	"
Methionine	1.80	1.60	1.51	0.62	"
Lysine	4.27	4.20	4.06	3.06	"
topsfeed	37.50	25.00	12.50	Nil	100




Pauletia seeds	12.50	25.00	37.50	50.00	Nil
Total	100kg	100.03kg	100kg	100.23kg	100kg

Falaye., 2014

Table 2: Level of Ant-nutritional factors in the raw and boiled seeds of *P. monandra*.

Components	T1 (Raw)	T2 (10 mins)	T3 (20 mins)	T4 (30 mins)	T5 (40 mins)	T6 (60 mins)
Phytate (mg/100g)	3.84 ^a ±0.4808	1.28 ^c ±0.3252	1.28 ^c ±0.0282	0.84 ^c ±0.0232	0.64 ^c ±0.0848	0.60 ^d ±0.1414
Hydrogen cyanide (mg/100g)	3.48 ^a ±0.2262	2.376 ^b ±0.0084	0.82 ^c ±0.0056	0.41 ^d ±0.0282	0.400 ^d ±0.0141	0.26 ^e ±0.0169
Tannin(%)	6.03 ^a ±0.0636	2.02 ^b ±0.2404	1.84 ^c ±0.0650	1.06 ^e ±0.0848	0.82 ^e ±0.0424	0.68 ^e ±0.2616
Saponin(%)	4.02 ^a ±0.4834	4.02 ^a ±0.0282	4.00 ^a ±0.9333	2.08 ^b ±0.0989	1.68 ^b ±0.0714	1.22 ^c ±0.1979

Means with the same superscripts across the row are not significantly different ($P>0.05$) Value = means + S.E



Experimental System and Feeding Trial

The concrete out door pond of (1.3x0.9x1m), were thoroughly washed and scrubbed with agricultural lime (CaCO_3) which served as disinfectant. The ponds were then rinsed and drained after 24 hours to clear the ponds of the lime. Thereafter each Pond was filled with 150litres of water from reserved tank in the garden, with sources from Baguwie reservoir water. With a labeled on each feeds assigned to experimental diet. The fifteen ponds were each stock with twenty fingerling of *C. gariepinus* selected randomly. The water in each pond was changed fortnightly to remove the faeces and left over food. The experimental fish *C. gariepinus* fingerlings were collected from the Genetic Improvement Laboratory, Funtua hactary units Nigeria. They were acclimatized for six days before the takeoff of the experiment, this was necessary to enable the fingerlings empty their stomach content and to force them to adjust to the new diet.



Construction of Pond



Filling of pond



Washing the pond with Lime



fertilization



Testing stage of the pond



Testing stage of the pond

Experimental Procedure

The experimental fish were randomly distributed at a stocking density of 20 fingerlings per tanks in triplicates. They were fed at 5% body weight twice daily morning and evening at equal ration. Sampling was done weekly using a sensitive electronic balance (Soehnle - 2000g Model) to determine the average weight of the fish and adjust the feed accordingly.



Acclimatization



Counting Stages



Introduction to pond



Feeding regime



Feeding regime



Growth stage

The study was conducted for 54 days. All analyses for proximate composition including the carcass composition before and after the experiment were determined according to the methods of (Association for Official Anyalytical Chemist) AOAC (2020). Water temperature was monitored daily



with a standardized mercury thermometer while dissolved oxygen and pH were determined using Digital DO meter and Jenway Automatic pH meter (Hannan HI98129) respectively.

Measurement of growth parameters.

Food conversion ratio (FCR), Specific growth rate (SGR) and Percentage survival rate were Determined as follows (After Fagbenro *et al.*, 1991).

$$FCR = \frac{\text{Dry Weight Feed Supplied}}{\text{Total Weight gain by fish (g)}}$$

$$SGR (\%) = 100 \frac{(\text{Log}^6 \text{Wt} - \text{Log}^6 \text{WO})}{T \text{ (days)}}$$

$$\text{Average Daily Wt gain} = \frac{\text{Wt} - \text{Wo}}{T \text{ (days)}}$$

$$S (\%) = \frac{N_t \times 100}{N_o}$$

where

Wt = Final body weight of fish in grams at the end of the experiment,

WO = Initial body weight of fish in grams at the beginning of experiments.

Log⁶ = Natural Logarithm of both final and initial body weight of fish in grams.

T = Duration (time) of the feeding trial in days.

No = Number of fingerlings alive at the end of experiment

S = % survival

Gross Protein Value (GPV): This is commonly used biological method for evaluating proteins. This was determined using Devendra (1988) method as.

GPV = A/AO where

A = $\frac{\text{g increase in weight gain}}{\text{g test protein}}$

AO = $\frac{\text{g increase, in weight gain}}{\text{g test protein}}$

Protein Intake (PI)

This was determined following Sveier *et al.*, (2000) method using the formula: PI = Total feed intake x % crude protein in the diet.

Protein efficiency ratio (PER)

This index uses growth as a measure of nutritive value of dietary protein. It was determined using Wilson (1989) as

$$PER = \frac{\text{Mean weight gain (g)}}{\text{Mean protein intake (mg)}}$$

Productive protein value (PPV)

This expresses the percentage of ingested protein that is retained by disposition in the carcass. It is usually calculated by the carcass analyses method of Miller and Bender (1955). When no correction for endogenous nitrogen losses is made the results are expressed as apparent net protein utilization (ANPU)



$$PPV (\%) = \frac{(P2-PI) \times 100}{\text{Total protein consumed.}}$$

Where PI is the protein in fish carcass (g) at the beginning of the experimental and P2 is the protein in fish carcass (g) at the end of the experiment.


Sampling Monitoring during the Experimental Fish; The length and weight of each fingerlings in each tank was measured at the commencement of the experiment. Subsequently, ten fingerlings were taken randomly from each tank once two week and weighed the growth rate. The sampling exercises were carried out in the morning before feeding the fish. Any dead fish is quickly removed and recorded to survival rate.

Monitoring of Water Quality: Water temperature record was taken daily before feeding (7.00-8.00) with Hanna instrument and water transparency secchi-disc thermometer. Dissolved oxygen and pH were determined using the methods described by Auta (1993)

Temperature; Temperatures readings were taken with HANNA Instrument HI 98129 dipping the mercury bulk in water for two minutes and then the value was recorded and reading on the calibrated glass stem, this was carried out weekly.

PH Hydrogen Ion Concentration ; pH determination was done by using HANNA Probe Meter Instrument HI -98129 by lowering the mercury bulk in water for two minutes and the value was recorded and reading on the calibrated glass stem, this was carried out bi-weekly.

Dissolved Oxygen (DO) ; This was determined be-weekly by using “Winkler” test-kit (Aqua Merck) instrument were used to determine the(DO) by lowering the top into the body of water and the reading were taken and Secchi disc were used for measure transparence of water before change it, show in the figure below



The following instruments were used during the experimental period for measuring; water dissolved oxygen, temperature, hydrogen ion consecration, water transparency and water acidity and alkanality.





Economic Analysis of Experimental Feeds

Weight gained in eight weeks of the experimental fish (*C.gariepinus fingerlings*) revealed that diet 1 (20.09g) was higher than all other diets, with diet 5 gaining an average weight of (18.60g) The least weight gained was observed in diet 4 (8.80g). Fish fed diets 2 and 3 gained average weights of (15.02g) and (11.02g) respectively. Fish fed diet 1 consumed the feed (36.44g), while those in diet 4 consumed the least (9.80g). The cost of diet 1 per kilograms was (N133.65), diet 2 (N117.16/kg), diet 3 (N99.61/kg), diet 4 (N87.55/kg) and diet 5 (N170.00/kg), and during the eight weeks period, the cost of feed consumed was N4.87 for diet 1, N4.00 for diet 2, N1.40 for diet 3, N 0.77 for diet 4 and N4.27 for diet 5. If a 24 weeks/6 months feeding period is considered, it will take about (N14.61) to feed and produce one fish with diet 1, (N12.00) for diet 2, (N4.17) for diet 3, (N2.31) for diet 4 and (N14.16) for diet 5. If one kilograms of fish is desired, it will cost about (N233.00) for diet 1 in 6 months, (N266.31) for diet 2 in 9 months, (N127.04) for diet 3 in 12 months, (N87.50) for diet 4 in 15 months and (N254.00) for diet 5 in 6 months. The projected profit expected at harvest time was N267.00 for diet 1, N233.69 for diet 2, N12.96 for diet 3, N412.50 for diet 4 and N246.00 for diet 5

Economic Analysis of Experimental Feeds

	Diet1	Diet2	Diet3	Diet4	Diet5
Weight gain(g)	20.09	15.02	11.02	8.80	18.60
Feed consumed (g)	36.44	34.15	14.00	9.80	27.74
Cost of feed/kg (n)	133.65	117.16	99.61	78.55	170
Cost of feed in 8weeks (n)	4.87	4.00	1.40	0.77	4.27
In 24 weeks (n)	14.61	12.00	4.17	2.31	14.16
Attain 1 kg (n)	233.00	266.31	127.04	87.50	254.00
Time (month)	6	9	12	15	6
Expected profit (n)	267.00	233.69	372.96	412.50	246.00

Linear Programing Showing Least Cost Formulation for Experimental Diets

Showed the possible combination of all the various amounts of each ingredient (g) and cost (N) needed in each diet at varying proportions in order to achieve 49% crude protein value of the feed at the most economical cost, gave the impetus for the calculated analysis of the experimental diets its indicates the amount of energy, crude protein, crude fibre, ether extract, potassium, magnesium, cysteine, lysine that was expected to be derived from each formulated diet after utilization by the fish. The calculated analysis also shows the cost of each diet per kilogram, with top fed as the most expensive (N170/kg) and the least D4 (N78.55). Diets 1, 2, 3 had cost values of N133.65, N117.16, and N99.61 respectively. The economy of weight gain showed that diet 1 was the most economical (N7.09/g) while diet 4, (N1.72/g) was the least economically.

Statistical Analysis

Data collected was subjected to analysis of variance (ANOVA) and the means from the various treatments were compared for significant differences ($p < 0.05$) using Duncan Multiple Range Test (DMRT) to rank the means, System Analytical Statistical (SAS) computer package version nine(9) was used for the analysis..

Anova for proximate composition of diets used for the experiment

Source of variation	Ss	df	ms	f	p-value	F crit
Between Group	29655.85	5	5931.17	1513.46	3.6E-29	2.620654
Within Group	94.05472	24	3.918947			
Total	29749.9	29				



Anova for growth performances and feeds utilization of *C. gariepinus* fingerlings

Source of variation	Ss	df	ms	f	p-value	F crit
Between Group	29655.85	5	5931.17	1513.46	3.6E-29	2.620654
Within Group	94.05472	24	3.918947			
Total	29749.9	29				

Table 3 Growth Performance of *C.gariepinus* fingerlines Fed Experimental Diets

Growth Parameter	Diets				
	1	2	3	4	5(control
Mean Initial body weight (g)	2.06	2.10	1.90	2.10	1.95
Mean final body weight (g)	20.90 ^a	15.02 ^b	11.02 ^b	8.80 ^c	18.60 ^a
Mean body weight gain (g)	18.94 ^a	12.92 ^b	9.12 ^b	6.70 ^c	16.80 ^a
Mean body weight gain (g)	914.56	615.24	480.00	319.05	862.05
Mean initial standard length (cm)	11.42 ^a	9.94 ^a	8.22 ^b	8.10 ^b	15.01 ^a
Mean standard length gain (cm)	5.86 ^a	3.84 ^a	2.32 ^b	2.10 ^b	8.99 ^a
Mean standard length gain	105.40	62.95	39.32	35.00	149.32
Specific growth rate (SGR)	4.14	3.51	3.14	2.56	4.03
Feed conversion ratio(FCR) %	51.70 ^d	37.83 ^c	65.14 ^b	83.75 ^a	60.60 ^c
Protein efficiency ratio (PER)	1.04 ^a	0.71 ^a	1.31 ^a	1.68 ^a	1.23 ^a
Net protein utilization (NPU) %	90.48	87.87	225.11	341.40	125.71
Conditional factor (K)	1.34	1.53	1.63	1.72	0.55

Means with the same superscripts across the rows are not significantly different ($P>0.05$)




Figure 1 Bi-weekly increases in body weight of *C. gariepinus*




Figure.11 Bi – weekly increase in standard length gain of *C. gariepinus* fingerlings

Discussion

In the present investigation, all the experimental diets were accepted by *C. gariepinus* fingerlings, indicating that the levels of incorporation of *P.monandra* meals did not affect the palatability of the diets. This portrays higher acceptability and utilization of the diets with additional advantage of being cheaper sources. This observation was supported by the work of Fagbenro, (2021) and Sotolu and Faturoti, (2017). These workers reported that reduction in anti-nutritional by different processing techniques resulted in better palatability and acceptability feed.

It was observed in this study that as temperature increased, pH increased and the DO value decreased. These parameters are within the range of culturing *C. gareipinus fingerlings*. Auta (1993) reported Temperature range of 25-30°C, Dissolved oxygen of 5mg/l and a pH range of 6.7-9.0 in a similar experiment. It is also important to note that the period of higher growth, both in weight and length, in this study, coincided with the period of higher temperature. (September to October, 2025) coincided with the warm /hot season in Kano. One of the factors affecting metabolic rate of fish is temperature. This parameter is directly affects the rate of metabolic activity of fish (Fakunle *et al.*, 2013).

Weight gain and standard length increases are known to be the most important indices for measuring fish responses to experimental diets and very liable indicators of growth (Balogun *et al.*, 2008, Tunde, *et al.*, (2018)). The study reveals that the specific growth rate per day of the fish was higher at diet 1. This might be an indication that the nutrients were best converted to flesh by the fish fed, at diet 4 level reduced the growth rate and feed utilization, which might be due to high fiber content in the diets. Contrary to observation made by Fafioye *et al.*, (2015). Balogun *et al.*, (2008), in other experimental diets containing processed *Delonix regia* seeds meal affected the growth *Oreochromis niloticus* at 12.5% inclusion did not affect growth. However, at high levels of inclusion, 25% or more, the growth of *O. niloticus* was high growth obtained Abegunde, and Odedire, (2015). The high values of performance indices (SGR and FCR) particularly in diet 1, 2, and 5 may be attributed to the high



protein content of the diets (approximately 47% and above) which is within the recommended range of dietary requirements for proper growth of *C. gariepinus*

The general robustness of the fish fed *P. monandra* meals at different level as expressed by the condition factor (K) shows that all fish were in better condition at the end of the experiment than at then gaining revealing that accumulation of real body weight and not fat is as a result of good protein utilization the outcome was in line with Oladapo et al, (2018), reporting on practical manual of *C. gariepinus* culture fish in Netherlands at different ranged of temperature a bit higher than a set temperature at end of the experiment, but contrary to observation made by Olasunkami, (2013) on growth performance and feed utilization of *Oreochromis niloticus* and *C. gariepinus* fingerlings fed on treated levels of *Delonix regia* seed meal that condition was no favours at the end of the experiment. The fish carcass composition shows that the crude protein content in all fish fed the experimental diets was higher than in fish before the experiment. This position is an indication that protein synthesis was achieved. Weight gain has been attributed to a clear indication of protein synthesis, increased tissue production and real growth which is in line with the reports of Olasunkanmi et al, (2013).

The combination diet, Diet 1 (topfeeds + *P. monandra* seeds) gave higher growth values over topfeeds alone and *P. monandra* diet alone, this portrays higher acceptability and utilization of the combination diet. An additional advantage of this combined diet is that it was also cheaper than the control diet (topfeeds alone). Similar cheaper combinations were reported by Hassan, Modu and Mohammed, (2017). It is also important to note that the period of higher growth, both in weight and length, in this study, coincided with the period of higher temperature. The period of the year when this experiment was conducted (November to December 2008) coincided with the cold harmattan season in Kano state, Nigeria. Coincidentally, there was increase in temperature from 23.10-24.20C at the last two weeks of the research (6"-8" week). This may have contributed to the high growth (length and weight) during the aforementioned period. One of the important factors affecting metabolic rate of fish is temperature. This parameter is closely linked and directly affects the rate of metabolic activity of fish (Olasunkanmi, 2012).

The relative cost of feed per unit weight gain or the economy of weight gain by the fish has proven that using top feed or *P. monandra* seed meal alone to supplement compounded diets would not be economical to fish growth. However, the combination of both (top feeds + *P. monandra* seed meal) would be more economical for weight gain and growth of fish. Therefore investment in low cost of feeding that translates to better growth rate and improved profit is advocated at this stage of global economic meltdown and high rate of exchange of the Nigerian Naira to Dollar. This study showed that *P. maonandra* meal, which is almost costless, could be used to partially replace the very expensive fishmeal in the diet of *C.gareipinus fingerlings* at regulated inclusion levels. This will no doubt reduce cost of production and thereby further boost aquaculture development in Nigeria.

References

- Abdullahi, S.A (2009). Determination of Protein Qualities of Three Species of Fresh Water Fishes, *Gynarchus niloticus*, *Heterotis niloticus* and *Latis niloticus* Using Chemical indices. *International Journal of Applied Biological Research*,. 1; 34-39.
- Abegunde, T. O. and Odedire, J. A. (2015). Nutritional assessment of leaves and seeds of some horticultural trees and shrubs as feed for ruminants in Southwest Nigeria. *Tropical. Animal. Production Investment*. <http://animalsci.agric.ui.edu.ng/TAPI/index.php/tapijournal/index>. University of Ibadan, 18 (1): 12-17.--
- Adewolu, M. F. (2008). Potential of Sweet Potato (*Ipomoea batatas*) Leaf Meal as Dietary Ingredient for *Tilapia zilia* Fingerlings, *Pakistan Journal of Nutrition*, 7 (3); 444- 449 Asian Network for Scientific Information
- Agbugui, M., Oniye, S.J., Auta, J. (2010). Effects of Processing on The Mineral Content, Proximate Composition and Phytochemical Factors of the Seeds of *Bauhinia Monandra* (KURZ). *Bayero Journal. Pure Applied. Sciences*. 2010 3(1): 23 – 25



- Anhwange, B.A, Ajibola, V. O., and Oniye, S. J. (2005). Nutritional Potential of the Seeds of *Bauhinia monandra* (Linn). *Journal of Food Technology*, 3(2):204-208.
- Anwa, E.P., Auta, J., Abdullahi, S.A and Bolorunduro, P.I. (2017). Effects of processing on seeds of *Albizia lebbek*: Proximate Analysis and Phytochemical Screening. *Research Journal of Biological Sciences* 2 (1): 41-44
- AOAC (2001). Official Methods of Analysis of the Association of official analytical chemist. .11.Washington, USA, 1234pp \
- Association of Official Analytical Chemists (2020). Official methods of chemical analysis 25th Edition, Washington D.C., USA.
- Auta, J. (1993). Water Quality Management in Fish Pond, In; Oniye, S. J, Bolurunduro, P.I and Auta J. (Ed) *Proceedings of National Workshop on Fisheries Extension Delivery*. NAERLS, A.B.U Zaria. .1-20.
- Balogun, J.K., A.S Abdullahi., J.Auta and O.P. Ogunlade, (2008). Feed Conversion, Protein Efficiency, Digestibility and Growth Performance of *Oreochromis niloticus* fed *Delonix regia* Seed Meal. 2008 In: *Proceedings of the National Conference of Fishers Society of Nigeria* (FISON). FISON, Lagos, Nigeria. 823-829 .
- Burke, R.L (2019). *An Economics Analysis of Producing Pond Raised Catfish for Food on Mississippi* Bull. No 870 Mississippi Agriculture and Forestry Experimentation Station, Mississippi State Ms. 25p.
- Fakunle, J. O., Alatise, S. P., Effiong, B. N. and Tihamiyu, K. (2013). Effects of Replacing Soyabeans Meal with Graded Levels of Effects of Replacing Soyabeans Meal with Graded Levels of Boiled *Jatropha* Kernel Meal in Diets of *Clarias gariepinus* Fingerlings. *Bulletin of Environmental, Pharmacological and Life Sciences*, 2 (9): 112-117.
- FAO, (2012). *Analysis of Feeds and Fertilizers for Sustainable Aquaculture Development* in Vietnam in Hassan, M.R. Hecht, T.; De silver Tacon A.G.J (Ed) study and technical paper no1497 Rome FAO2007. 331 – 357
- FAO, (2020). *Analysis of Feeds and Fertilizers for Sustainable Aquaculture Development* in Vietnam in Hassan, M.R. Hecht, T.; De silver Tacon A.G.J (Ed) study and technical paper no 497 Rome FAO2007. 331 – 357
- Ferouz, Y.A., Kurt, A.R. and Kasiuviswanathan, M. (2013). Alternative Protein Sources for Aquaculture feeds. *J. Aquaculture Feed Sciences Nutrition*. 4(1):1-26.
- Hassan, M., Modu, M. and Mohammed, A. A. (2017). Effects of Water Melon (*Citrullus 116 lanatus*) on Growth Performance of *Oreochromis niloticus* Fingerlings as a Replacement for Soybean Meal. African Research Corps *International Journal of Agro-business and Agricultural Research*. 4(1):1-10
- Mamudu, .S.T, Musilimat.T. and Wuild A.A. (2018) Growth Response, Feed Conversion Rate and Cost Benefits of *C. gariepinus* Juveniles fed on processed Cassava leaf meal in outdoor Concrete Ponds, *International Journal of Innovation, Nutrition and Sustainable Agriculture*.6,4,25-35
- Olaniyi, C.O.; Aderinola, O.A.; Rafiu, T.A.; Akande, T.O. and Okunola, D.O. (2009b). Growth Performance and Nutrient Utilization of *Clarias gariepinus* fed Locust bean seed meal. *Proceedings of 14 Annual Conference of Animal Science Association of Nigeria (ASAN)* September 14 -17 held at LAUTECH, Ogbomosho, Nigeria. (Eds) Akinlade, J.A.; Olayeni, T.B; Rafiu, T.A.; Akinwumi, A.O; Aderinola, O.A; Ojebiyi, O.O. and Odunsi, A.A. 550-552.
- Olapade A. A., Ajayi O. A. and Olaniyi M. B. (2018). Proximate, Phytochemical and Mineral Compositions of Roasted Seeds of Coffee Senna (*Senna occidentalis* Linn). *Journal of Food Science and Technology*, 19 (1):51p.
- Olasunkanmi, J. B. (2013). Nutrient utilisation and growth performance of *Clarias gariepinus* fed differently processed *Mucuna utilis* meals as a replacement for soyabean based diet. PhD. Thesis, University of Ibadan, Ibadan, Nigeria. 209p.



- Salgueiro, A. C. F.(2022) The influence of Bauhinia forficata Link subsp. pruinosa tea on lipid peroxidation and non-protein SH groups in human erythrocytes exposed to high glucose concentrations. *Journal of Ethnopharmacology*, 148, 1, 81-87.
- Sotolu, A. O. and Faturoti, E. O. (2017).Digestibility and Nutritional Values of Differently Processed *Leucaena leucocephala* (Lam De Wit) Seed Meals in the Diet of African Catfish (*Clarias gariepinus*). *Middle East Journal of Scientific Research*, 3 (4): 190
- Teixeira, D. C., Farias D. F., Urano-Carvalho, A. F., Arantes, M. R., Oliveira, J.T., Sousa, D.O., Pereira, M. L., Oliveira, H. D., Andrade Neto, M., and Vasconcelos, I. M. (2013) Chemical composition, nutritive value, and toxicological evaluation of Bauhinia cheilantha Seeds: A legume from semiarid regions widely used in folk medicine. *Bio-Medical Research International*, 2013: 578781.
- Tunde, A. O.,Oluwagbemiga, O. S.,Babatunde, A.and Oluseyi,O. O.(2018).The Growth Performance of African Catfish (*Clarias gariepinus*) Fed Commercially Prepared Imported Fish Feeds. *International Journal of Agricultural Economics*.1 (3): 57-61.
- Umaru, J., Auta, J., Oniye, S. and Bolorunduro, P. (2016).“Growth and economic performance of African catfish, *Clarias gariepinus* (Burchell) Juveniles to imported and local feeds in floating bamboo cages”. *International Journal of Fisheries and Aquatic Studies*, 4 (2): 221-226.

Acknowledgments: The researchers would like to gratefully thank the Tertiary Education Trust Fund (TETFUND) for sponsoring the research.